ľ.

|st

"!]

iji iji

:=4

Jan Matthijs JETTEN et al. - Docket No. BO 42503

especially 2, 2, 6, 6-tetramethylpiperidin-1-oxyl (TEMPO).

- --21. (New) A process according to Claim 19, wherein the enzyme capable of oxidation is an oxidoreductase.
- --22. (New) A process according to Claim 21, wherein the enzyme is a peroxidase, especially horse radish, soy-bean, lignin peroxidase or myelo- or lacto-peroxidase, and the oxidizing agent is hydrogen peroxide.
- $^{--23}$. (New) A process according to Claim 21, wherein the enzyme is a polyphenol oxidase or a laccase and the oxidizing agent is oxygen.
- --24. (New) A process according to Claim 19, wherein the enzyme is a hydrolase, especially phytase or lipase, in the presence of a metal compound.
- --25. (New) A process according to Claim 19, wherein the primary alcohol is comprised in a carbohydrate.
- $^{--26.}$ (New) A process according to Claim 25, wherein the carbohydrate is an $\alpha\mbox{-glucan}$ or fructan or a derivative thereof.
- --27. (New) A process according to Claim 25, wherein a carbonyl-containing carbohydrate containing at least 1 cyclic monosaccharide chain group carrying a carbaldehyde group per 25 monosaccharide units and per average molecule is produced.
- --28. (New) A process according to Claim 25, wherein the carbohydrate is a hydroxyalkylated carbohydrate or a glycoside or a glyconic acid.

Jan Matthijs JETTEN et al. - Docket No. BO 42503

- --29. (New) A process according to Claims 19, wherein the primary alcohol is comprised in a steroid compound.
- --30. (New) A process according to Claim 19, wherein the primary alcohol is comprised in textile fibers.
- --31. (New) An oxidized carbohydrate, the carbohydrate being selected from disaccharides, oligosaccharides and polysaccharides of the glucan, mannan, fructan, and chitin types galactan, and carbohydrate glycoside, containing at least 1 cyclic monosaccharide chain group carrying a carbaldehyde group per 25 monosaccharide units and per average molecule, or a chemical derivative thereof.
- --32. (New) An oxidized carbohydrate according to Claim 31, containing at least 5 monosaccharide units per average molecule.
- --33. (New) An oxidized carbohydrate according to Claim 31, which contains 1 to 50 cyclic monosaccharide chain group carrying a carbaldehyde group per 50 monosaccharide units and per average molecule.
- --34. (New) A carbohydrate derivative according to Claim 31, in which derivative at least a part of the carbaldehyde groups has been converted to a group with the formula -CH=N-R or -CH₂-NHR, wherein R is hydrogen, hydroxyl, amino, or a group R^1 , OR^1 or NHR^1 , in which R^1 is C_1 - C_{20} alkyl, C_1 - C_{20} acyl, a carbohydrate residue, or group coupled with or